

1. Singer AJ, Clark RA. Cutaneous wound healing. *N Engl J Med*. Sep 2 1999;341(10):738-46. doi:10.1056/NEJM199909023411006
2. Son D, Harijan A. Overview of surgical scar prevention and management. *J Korean Med Sci*. Jun 2014;29(6):751-7. doi:10.3346/jkms.2014.29.6.751
3. Finnsen KW, McLean S, Di Guglielmo GM, Philip A. Dynamics of Transforming Growth Factor Beta Signaling in Wound Healing and Scarring. *Adv Wound Care (New Rochelle)*. Jun 2013;2(5):195-214. doi:10.1089/wound.2013.0429
4. Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol*. May 2002;3(5):349-63. doi:10.1038/nrm809
5. Alexander Nast GG, Kerstin Lorenz, Hans-Robert Metelmann, Uwe Paasch, Vratislav Strnad, Michael Weidmann, Ricardo Niklas Werner, Jürgen Bauerschmitz. S2k guidelines for the therapy of pathological scars (hypertrophic scars and keloids) – Update 2020. *AWMF Registry No: 013-030, 2020*. 04 October 2020;
6. Redd MJ, Cooper L, Wood W, Stramer B, Martin P. Wound healing and inflammation: embryos reveal the way to perfect repair. *Philos Trans R Soc Lond B Biol Sci*. May 29 2004;359(1445):777-84. doi:10.1098/rstb.2004.1466
7. Moon J, Yoon JY, Yang JH, Kwon HH, Min S, Suh DH. Atrophic acne scar: a process from altered metabolism of elastic fibres and collagen fibres based on transforming growth factor-beta1 signalling. *Br J Dermatol*. Dec 2019;181(6):1226-1237. doi:10.1111/bjd.17851
8. Vorstenbosch J, Gallant-Behm C, Trzeciak A, Roy S, Mustoe T, Philip A. Transgenic mice overexpressing CD109 in the epidermis display decreased inflammation and granulation tissue and improved collagen architecture during wound healing. *Wound Repair Regen*. Mar-Apr 2013;21(2):235-46. doi:10.1111/wrr.12023
9. Gauglitz GG, Korting HC, Pavicic T, Ruzicka T, Jeschke MG. Hypertrophic scarring and keloids: pathomechanisms and current and emerging treatment strategies. *Mol Med*. Jan-Feb 2011;17(1-2):113-25. doi:10.2119/molmed.2009.00153
10. Scott PG, Dodd CM, Tredget EE, Ghahary A, Rahemtulla F. Immunohistochemical localization of the proteoglycans decorin, biglycan and versican and transforming growth factor-beta in human post-burn hypertrophic and mature scars. *Histopathology*. May 1995;26(5):423-31. doi:10.1111/j.1365-2559.1995.tb00249.x
11. Xie JL, Qi SH, Pan S, et al. Expression of Smad protein by normal skin fibroblasts and hypertrophic scar fibroblasts in response to transforming growth factor beta1. *Dermatol Surg*. Sep 2008;34(9):1216-24; discussion 1224-5. doi:10.1111/j.1524-4725.2008.34261.x
12. Grieb G, Steffens G, Pallua N, Bernhagen J, Bucala R. Circulating fibrocytes--biology and mechanisms in wound healing and scar formation. *Int Rev Cell Mol Biol*. 2011;291:1-19. doi:10.1016/B978-0-12-386035-4.00001-X
13. Wang JF, Jiao H, Stewart TL, Shankowsky HA, Scott PG, Tredget EE. Fibrocytes from burn patients regulate the activities of fibroblasts. *Wound Repair Regen*. Jan-Feb 2007;15(1):113-21. doi:10.1111/j.1524-475X.2006.00192.x
14. Sarrazy V, Billet F, Micallef L, Coulomb B, Desmouliere A. Mechanisms of pathological scarring: role of myofibroblasts and current developments. *Wound Repair Regen*. Sep 2011;19 Suppl 1:s10-5. doi:10.1111/j.1524-475X.2011.00708.x
15. Dulauroy S, Di Carlo SE, Langa F, Eberl G, Peduto L. Lineage tracing and genetic ablation of ADAM12(+) perivascular cells identify a major source of profibrotic cells during acute tissue injury. *Nat Med*. Aug 2012;18(8):1262-70. doi:10.1038/nm.2848

16. Phan TT, Lim IJ, Aalami O, et al. Smad3 signalling plays an important role in keloid pathogenesis via epithelial-mesenchymal interactions. *J Pathol*. Oct 2005;207(2):232-42. doi:10.1002/path.1826
17. Monstrey S, Middelkoop E, Vranckx JJ, et al. Updated scar management practical guidelines: non-invasive and invasive measures. *J Plast Reconstr Aesthet Surg*. Aug 2014;67(8):1017-25. doi:10.1016/j.bjps.2014.04.011
18. Young VL, Hutchison J. Insights into patient and clinician concerns about scar appearance: semiquantitative structured surveys. *Plast Reconstr Surg*. Jul 2009;124(1):256-265. doi:10.1097/PRS.0b013e3181a80747
19. Chen MA, Davidson TM. Scar management: prevention and treatment strategies. *Curr Opin Otolaryngol Head Neck Surg*. Aug 2005;13(4):242-7. doi:10.1097/01.moo.0000170525.74264.f8
20. Mustoe TA, Cooter RD, Gold MH, et al. International clinical recommendations on scar management. *Plast Reconstr Surg*. Aug 2002;110(2):560-71. doi:10.1097/00006534-200208000-00031
21. Niessen FB, Spauwen PH, Schalkwijk J, Kon M. On the nature of hypertrophic scars and keloids: a review. *Plast Reconstr Surg*. Oct 1999;104(5):1435-58. doi:10.1097/00006534-199910000-00031
22. Wittenberg GP, Fabian BG, Bogomilsky JL, et al. Prospective, single-blind, randomized, controlled study to assess the efficacy of the 585-nm flashlamp-pumped pulsed-dye laser and silicone gel sheeting in hypertrophic scar treatment. *Arch Dermatol*. Sep 1999;135(9):1049-55. doi:10.1001/archderm.135.9.1049
23. Suetake T, Sasai S, Zhen YX, Ohi T, Tagami H. Functional analyses of the stratum corneum in scars. Sequential studies after injury and comparison among keloids, hypertrophic scars, and atrophic scars. *Arch Dermatol*. Dec 1996;132(12):1453-8.
24. Poetschke J, Gauglitz GG. Treatment of Immature Scars: Evidence-Based Techniques and Treatments. In: Teot L, Mustoe TA, Middelkoop E, Gauglitz GG, eds. *Textbook on Scar Management: State of the Art Management and Emerging Technologies*. 2020:193-201.
25. Cho YS, Jeon JH, Hong A, et al. The effect of burn rehabilitation massage therapy on hypertrophic scar after burn: a randomized controlled trial. *Burns*. Dec 2014;40(8):1513-20. doi:10.1016/j.burns.2014.02.005
26. Shin TM, Bordeaux JS. The role of massage in scar management: a literature review. *Dermatol Surg*. Mar 2012;38(3):414-23. doi:10.1111/j.1524-4725.2011.02201.x
27. Ngaage M, Agius M. The Psychology of Scars: A Mini-Review. *Psychiatr Danub*. Nov 2018;30(Suppl 7):633-638.
28. Manuskiatti W, Fitzpatrick RE. Treatment response of keloidal and hypertrophic sternotomy scars: comparison among intralesional corticosteroid, 5-fluorouracil, and 585-nm flashlamp-pumped pulsed-dye laser treatments. *Arch Dermatol*. Sep 2002;138(9):1149-55. doi:10.1001/archderm.138.9.1149
29. Wick G, Grundtman C, Mayerl C, et al. The immunology of fibrosis. *Annu Rev Immunol*. 2013;31:107-35. doi:10.1146/annurev-immunol-032712-095937
30. Coentro JQ, Pugliese E, Hanley G, Raghunath M, Zeugolis DI. Current and upcoming therapies to modulate skin scarring and fibrosis. *Adv Drug Deliv Rev*. Jun 2019;146:37-59. doi:10.1016/j.addr.2018.08.009
31. Oliveira GV, Hawkins HK, Chinkes D, et al. Hypertrophic versus non hypertrophic scars compared by immunohistochemistry and laser confocal microscopy: type I and III collagens. *Int Wound J*. Dec 2009;6(6):445-52. doi:10.1111/j.1742-481X.2009.00638.x

32. Berman B, Flores F. Recurrence rates of excised keloids treated with postoperative triamcinolone acetonide injections or interferon alfa-2b injections. *J Am Acad Dermatol*. Nov 1997;37(5 Pt 1):755-7. doi:10.1016/s0190-9622(97)70113-0
33. Berman B, Duncan MR. Short-term keloid treatment in vivo with human interferon alfa-2b results in a selective and persistent normalization of keloidal fibroblast collagen, glycosaminoglycan, and collagenase production in vitro. *J Am Acad Dermatol*. Oct 1989;21(4 Pt 1):694-702. doi:10.1016/s0190-9622(89)70239-5
34. Nanda S, Reddy BS. Intralesional 5-fluorouracil as a treatment modality of keloids. *Dermatol Surg*. Jan 2004;30(1):54-6; discussion 56-7. doi:10.1111/j.1524-4725.2004.29382.x
35. Skochdopole A, Dibbs RP, Sarrami SM, Dempsey RF. Scar Revisions. *Semin Plast Surg*. May 2021;35(2):130-138. doi:10.1055/s-0041-1727291
36. Reish RG, Eriksson E. Scar treatments: preclinical and clinical studies. *J Am Coll Surg*. Apr 2008;206(4):719-30. doi:10.1016/j.jamcollsurg.2007.11.022
37. Zuber TJ, DeWitt DE. Earlobe keloids. *Am Fam Physician*. Jun 1994;49(8):1835-41.
38. Alster TS, Lupton JR. Lasers in dermatology. An overview of types and indications. *Am J Clin Dermatol*. 2001;2(5):291-303. doi:10.2165/00128071-200102050-00004
39. Anderson RR, Parrish JA. Selective photothermolysis: precise microsurgery by selective absorption of pulsed radiation. *Science*. Apr 29 1983;220(4596):524-7. doi:10.1126/science.6836297
40. Leszczynski R, da Silva CA, Pinto A, Kuczynski U, da Silva EM. Laser therapy for treating hypertrophic and keloid scars. *Cochrane Database Syst Rev*. Sep 26 2022;9(9):CD011642. doi:10.1002/14651858.CD011642.pub2
41. Nouri K, Vidulich K, Rivas MP. Lasers for scars: a review. *J Cosmet Dermatol*. Mar 2006;5(1):14-22. doi:10.1111/j.1473-2165.2006.00217.x
42. Ogawa R, Mitsuhashi K, Hyakusoku H, Miyashita T. Postoperative electron-beam irradiation therapy for keloids and hypertrophic scars: retrospective study of 147 cases followed for more than 18 months. *Plast Reconstr Surg*. Feb 2003;111(2):547-53; discussion 554-5. doi:10.1097/01.PRS.0000040466.55214.35
43. Atiyeh BS. Nonsurgical management of hypertrophic scars: evidence-based therapies, standard practices, and emerging methods. *Aesthetic Plast Surg*. Sep-Oct 2007;31(5):468-92; discussion 493-4. doi:10.1007/s00266-006-0253-y
44. Leventhal D, Furr M, Reiter D. Treatment of keloids and hypertrophic scars: a meta-analysis and review of the literature. *Arch Facial Plast Surg*. Nov-Dec 2006;8(6):362-8. doi:10.1001/archfaci.8.6.362
45. Rusciani L, Rossi G, Bono R. Use of cryotherapy in the treatment of keloids. *J Dermatol Surg Oncol*. Jun 1993;19(6):529-34. doi:10.1111/j.1524-4725.1993.tb00386.x
46. Har-Shai Y, Amar M, Sabo E. Intralesional cryotherapy for enhancing the involution of hypertrophic scars and keloids. *Plast Reconstr Surg*. May 2003;111(6):1841-52. doi:10.1097/01.PRS.0000056868.42679.05
47. Negenborn VL, Groen JW, Smit JM, Niessen FB, Mullender MG. The Use of Autologous Fat Grafting for Treatment of Scar Tissue and Scar-Related Conditions: A Systematic Review. *Plast Reconstr Surg*. Jan 2016;137(1):31e-43e. doi:10.1097/PRS.0000000000001850
48. Prantl L, Eigenberger A, Reinhard R, Siegmund A, Heumann K, Felthaus O. Cell-Enriched Lipotransfer (CELT) Improves Tissue Regeneration and Rejuvenation without Substantial Manipulation of the Adipose Tissue Graft. *Cells*. Oct 8 2022;11(19)doi:10.3390/cells11193159

49. Lenzi L, Santos J, Raduan Neto J, Fernandes CH, Faloppa F. The Patient and Observer Scar Assessment Scale: Translation for portuguese language, cultural adaptation, and validation. *Int Wound J*. Dec 2019;16(6):1513-1520. doi:10.1111/iwj.13228
50. Krastev TK, Schop SJ, Hommes J, Piatkowski A, van der Hulst R. Autologous fat transfer to treat fibrosis and scar-related conditions: A systematic review and meta-analysis. *J Plast Reconstr Aesthet Surg*. Nov 2020;73(11):2033-2048. doi:10.1016/j.bjps.2020.08.023
51. Bassetto F, Scarpa C, Vindigni V. Invasive Techniques in Scar Management: Fat Injections. In: Téot L, Mustoe TA, Middelkoop E, Gauglitz GG, eds. *Textbook on Scar Management: State of the Art Management and Emerging Technologies*. Springer International Publishing; 2020:333-342.
52. Antar SA, Ashour NA, Marawan ME, Al-Karmalawy AA. Fibrosis: Types, Effects, Markers, Mechanisms for Disease Progression, and Its Relation with Oxidative Stress, Immunity, and Inflammation. *Int J Mol Sci*. Feb 16 2023;24(4)doi:10.3390/ijms24044004